

Spontaneous Partial Breaking of $\mathcal{N} = 2$ Supersymmetry and the $U(N)$ Gauge Model ¹

K. Fujiwara*, H. Itoyama* and M. Sakaguchi[†]

*Department of Mathematics and Physics, Graduate School of Science, Osaka City University
3-3-138, Sugimoto, Sumiyoshi-ku, Osaka, 558-8585, Japan

[†]Okayama Institute for Quantum Physics, 1-9-1 Kyoyama, Okayama 700-0015, Japan

Abstract. We briefly review the properties of the $\mathcal{N} = 2$ $U(N)$ gauge model with/without matters. On the vacua, $\mathcal{N} = 2$ supersymmetry and the gauge symmetry are spontaneously broken to $\mathcal{N} = 1$ and a product gauge group, respectively. The masses of the supermultiplets appearing on the $\mathcal{N} = 1$ vacua are given. We also discuss the relation to the matrix model.

Keywords: extended supersymmetry; spontaneous partial supersymmetry breaking

PACS: 11.30.Pb; 11.30.Qc; 11.25.Tq

The supercurrent algebra plays a key role in the partial breaking of global extended supersymmetries

$$\{\bar{Q}_{\alpha}^J, \mathcal{S}_{\alpha I}^m(x)\} = 2(\sigma^n)_{\alpha\dot{\alpha}} \delta_I^J T_n^m(x) + (\sigma^m)_{\alpha\dot{\alpha}} C_I^J \quad (1)$$

where $\mathcal{S}_{\alpha I}^m$ are extended supercurrents, T_n^m is the stress-energy tensor and C_I^J is a field independent constant matrix, which is permitted by the constraint for the Jacobi identities [1]. The new term does not modify the supersymmetry algebra on the fields.

Besides active researches on the *non-linear* realization of extended supersymmetry in the partially broken phase, Antoniadis, Partouche and Taylor (APT) [2] (see also [3]) gave a model in which *linearly* realized $\mathcal{N} = 2$ supersymmetry is partially broken to $\mathcal{N} = 1$ spontaneously. APT model is $\mathcal{N} = 2$ supersymmetric, self-interacting $U(1)$ model composed of one (or several) abelian constrained $\mathcal{N} = 2$ vector multiplet(s) with electric & magnetic Fayet-Iliopoulos (FI) terms. In [4], we have generalized this model to the $U(N)$ gauge model and shown that the $\mathcal{N} = 2$ supersymmetry is partially broken to $\mathcal{N} = 1$ spontaneously. Further in [5], we have analyzed the vacua with broken gauge symmetry and revealed the $\mathcal{N} = 1$ supermultiplets on the vacua. The relation to the matrix model is discussed in [6].² In addition, a manifestly $\mathcal{N} = 2$ supersymmetric formulation of the model coupled with/without $\mathcal{N} = 2$ hypermultiplets was given in [9] by using unconstrained $\mathcal{N} = 2$ superfields on harmonic superspace [10]. These models contain the prepotential $\mathcal{F}$ as a prime ingredient. So, our model should be regarded as

¹ Talks given by HI at 14th International Conference on Supersymmetry and the Unification of Fundamental Interactions (SUSY06), Irvine, California, USA, 12-17 June 2006, and given by KF at MSJ-IHES Joint Workshop on Noncommutativity, Bures-sur-Yvette, France, 15-18 November 2006.

² This series of works [4, 5, 6] is based on $\mathcal{N} = 1$ superspace and construction of the most general $\mathcal{N} = 2$ Lagrangian based on special Kähler geometry [7] which was developed after tensor calculus [8].

a low-energy effective action for systems given by $\mathcal{N} = 2$ bare actions spontaneously broken to $\mathcal{N} = 1$.

We introduce $\mathcal{N} = 1$ superfields: chiral $\Phi(x^m, \theta) = \Phi^a t_a \ni (A, \psi, F)$ and vector $V(x^m, \theta) = V^a t_a \ni (v_m, \lambda, D)$, where t_a ($a = 0, \dots, N^2 - 1$) generate $u(N)$ algebra, $[t_a, t_b] = if_{ab}^c t_c$ and t_0 generates the overall $u(1)$. The model is composed of the Kähler term

$$\mathcal{L}_K = \int d^2\theta d^2\bar{\theta} K(\Phi^a, \Phi^{*a}), \quad K = \frac{i}{2}(\Phi^a \mathcal{F}_a^* - \Phi^{*a} \mathcal{F}_a), \quad (2)$$

where $\mathcal{F}_a = \frac{\partial \mathcal{F}}{\partial \Phi^a}$, the $U(N)$ gauging counterterm $\mathcal{L}_\Gamma$, the $U(N)$ gauge kinetic term

$$\mathcal{L}_{\mathcal{W}^2} = -\frac{i}{4} \int d^2\theta \mathcal{F}_{ab} \mathcal{W}^a \mathcal{W}^b + c.c. \quad \mathcal{W}_\alpha = -\frac{1}{4} \bar{D}\bar{D}e^{-V} D_\alpha e^V = \mathcal{W}_\alpha^a t_a, \quad (3)$$

a gauge invariant superpotential term and the FI D-term (e and m are real constants)

$$\mathcal{L}_W = \int d^2\theta W(\Phi) + c.c., \quad \mathcal{L}_D = \xi \int d^2\theta d^2\bar{\theta} V^0 = \sqrt{2}\xi D^0, \quad W = eA^0 + m\mathcal{F}_0. \quad (4)$$

In [4], it is shown that the action is invariant under $\mathfrak{R}$ -action which is composed of a discrete element of the $SU(2)$ R-symmetry and a sign flip of the FI parameter

$$R: \lambda_I^a \rightarrow \varepsilon^{IJ} \lambda_J^a \quad \& \quad R_\xi: \xi \rightarrow -\xi, \quad (5)$$

where $\lambda_I^a = (\frac{\lambda^a}{\psi^a})$, so that $S^{(+\xi)} \xrightarrow{R} S^{(-\xi)} \xrightarrow{R_\xi} S^{(+\xi)}$. We have made the sign of the FI parameter manifest. This ensures the $\mathcal{N} = 2$ supersymmetry of our action. In fact, acting $\mathfrak{R}$ on the first supersymmetry transformation $\delta_{\eta_1} S^{(+\xi)} = 0$, we have, $\delta_{\eta_1} S^{(+\xi)} = 0 \xrightarrow{R} R(\delta_{\eta_1}) S^{(-\xi)} = 0 \xrightarrow{R_\xi} \mathfrak{R}(\delta_{\eta_1}) S^{(+\xi)} = 0$, which implies that the resulting $\mathfrak{R}$ -invariant action is invariant under the second supersymmetry $\delta_{\eta_2} \equiv \mathfrak{R}(\delta_{\eta_1})$ as well. By applying the $\mathfrak{R}$ -action on the first supersymmetry transformation, we obtain the $\mathcal{N} = 2$ supersymmetry transformation of the fermion as

$$\delta \lambda_J^a = i(\tau \cdot \tilde{\mathbf{D}}^a)_J^K \eta_K + \dots, \quad \tilde{\mathbf{D}}^a = -\sqrt{2}g^{ab*} \partial_{b*} (\mathcal{E} A^{*0} + \mathcal{M} \mathcal{F}_0^*) \quad (6)$$

where $g_{ab} = \text{Im} \mathcal{F}_{ab}$ and τ are Pauli matrices. The rigid $SU(2)$ has been fixed by making $\mathcal{E}$ and $\mathcal{M}$ point to specific directions, $\mathcal{E} = (0, -e, \xi)$ and $\mathcal{M} = (0, -m, 0)$. Under the symplectic transformation, $\Omega = \begin{pmatrix} A^0 \\ \mathcal{E} \end{pmatrix} \rightarrow \Lambda \Omega$, $\Lambda \in Sp(2, \mathbb{R})$, $\begin{pmatrix} -\mathcal{M} \\ \mathcal{E} \end{pmatrix}$ changes to $\Lambda^{-1} \begin{pmatrix} -\mathcal{M} \\ \mathcal{E} \end{pmatrix}$. So the electric and magnetic charges are interchanged $(\mathcal{E}', \mathcal{M}') = (\mathcal{M}, -\mathcal{E})$ when $\Lambda = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. This explains the name of the electric and magnetic FI terms.

The vacua are specified by the scalar potential given by

$$\mathcal{V} = \frac{1}{8} g_{ab} \mathcal{P}^a \mathcal{P}^b + \frac{1}{2} g_{ab} \tilde{\mathbf{D}}^a \cdot \tilde{\mathbf{D}}^{*b}, \quad \mathcal{P}^a \equiv g^{ab} \mathcal{P}_b = -i f_{bc}^a A^{*b} A^c. \quad (7)$$

We find that for the case with $\mathcal{F} = \sum_{\ell=0} \frac{C_\ell}{\ell!} \text{Tr} \Phi^\ell$, the vacuum expectation values are determined in the eigenvalue basis by

$$\langle\langle \mathcal{F}_{\underline{ii}} \rangle\rangle = -2 \left(\frac{e}{m} \pm i \frac{\xi}{m} \right). \quad (8)$$

The vev of $\mathcal{V}$ is given by $\langle\langle\mathcal{V}\rangle\rangle = 2|m\xi|$. On the other hand, by applying supersymmetry transformation twice on the $U(1)_R$ charge conservation law [12], we can read off the constant matrix C_I^J in (1) as $C_I^J = +4m\xi(\tau_1)_I^J$. Thus half of supercurrents annihilates the vacuum.

Let $\langle\langle A \rangle\rangle$ be $\langle\langle A \rangle\rangle = \text{diag}(\overbrace{\lambda^{(1)}, \dots, \lambda^{(1)}}^{N_1}, \overbrace{\lambda^{(2)}, \dots, \lambda^{(2)}}^{N_2}, \dots)$ with $\sum_i N_i = N$ where $\lambda^{(k)}$ are complex roots of (8), then $U(N)$ is broken to $\prod_i U(N_i)$. Unbroken $\prod_i U(N_i)$ is generated by $t_a \in \{t_a | [t_a, \langle\langle A \rangle\rangle] = 0\}$, while broken ones by $t_\mu \in \{t_a | [t_a, \langle\langle A \rangle\rangle] \neq 0\}$. The $\mathcal{N} = 2$ supersymmetry is partially broken on the vacua since

$$\langle\langle \frac{1}{\sqrt{2}} \delta(\lambda^i \pm \psi^i) \rangle\rangle = \pm im \sqrt{\frac{2}{N}} (\eta_1 \mp \eta_2), \quad \langle\langle \frac{1}{\sqrt{2}} \delta(\lambda^i \mp \psi^i) \rangle\rangle = 0, \quad \langle\langle \delta \lambda_I^r \rangle\rangle = 0. \quad (9)$$

The mass spectrum is summarized as ($d_u \equiv \dim \prod_i U(N_i)$)

field	mass	label	# of polarization states
$v_m^\alpha, \frac{1}{\sqrt{2}}(\lambda^\alpha \pm \psi^\alpha)$	0	A	$2d_u$
$A^\alpha, \frac{1}{\sqrt{2}}(\lambda^\alpha \mp \psi^\alpha)$	$ m \langle\langle g^{\alpha\alpha} \mathcal{F}_{0\alpha\alpha} \rangle\rangle $	B	$2d_u$
$v_m^\mu + (TA^\mu)_2, \lambda_I^\mu$	$\frac{1}{\sqrt{2}} f_{\mu i}^\mu \lambda^{*i} $	C	$4(N^2 - d_u)$

(10)

$\frac{1}{\sqrt{2}}(\lambda^0 \pm \psi^0)$ is massless and thus is the Nambu-Goldstone (NG) fermion for the partial breaking of $\mathcal{N} = 2$ supersymmetry to $\mathcal{N} = 1$. Due to the $\mathcal{N} = 1$ supersymmetry on the vacua, the modes in (10) form $\mathcal{N} = 1$ multiplets: (A) massless $\mathcal{N} = 1$ vector multiples of $\text{spin}(\frac{1}{2}, 1)$, (B) massive $\mathcal{N} = 1$ chiral multiplets of $\text{spin}(0, \frac{1}{2})$ and (C) two massive $\mathcal{N} = 1$ vector multiplets of $\text{spin}(0, \frac{1}{2}, 1)$.

In [4], it was shown that the fermionic shift symmetry arises in a certain limit of our model, due to second supersymmetry, which is spontaneously broken. More precisely in [6], we study the decoupling limit of the NG fermion and the fermionic shift symmetry in $\mathcal{N} = 1$ $U(N)$ gauge model. The fermionic shift symmetry plays a key role in the proof of the Dijkgraaf-Vafa conjecture which asserts that non-perturbative quantities in $\mathcal{N} = 1$ supersymmetric gauge theory can be computed by a matrix model. The $\mathcal{N} = 1$ action was obtained by “softly” breaking of $\mathcal{N} = 2$ supersymmetry by adding the tree-level superpotential $\int d^2\theta \text{Tr} W(\Phi)$. Thanks to the fermionic shift symmetry, effective superpotential is written as $W_{\text{eff}} = \int d^2\chi \mathcal{F}_p$ with some function $\mathcal{F}_p$, which is related to the free energy of the matrix model [11].

We consider the $\mathcal{N} = 1$ $U(N)$ action realized on the vacua, in which the discrete $SU(2)$ R-symmetry is broken by the superpotential. How can we realize the situation in which the NG fermion is decoupled while the superpotential remains non-trivial? The answer is to consider the large limit of the electric and magnetic FI parameters (e, m, ξ). Parametrizing

$$(e, m, \xi) = (\Lambda e', \Lambda m', \Lambda \xi'), \quad \mathcal{F} = \text{tr} \left(c_0 \mathbf{1} + c_1 \Phi + \frac{c_2}{2} \Phi^2 \right) + \frac{1}{\Lambda} \sum_{\ell=3}^n \text{tr} \frac{c'_\ell}{\ell!} \Phi^\ell \quad (11)$$

and taking the limit $\Lambda \rightarrow \infty$, we derived the general $\mathcal{N} = 1$ action discussed in [11], in which the NG fermion is decoupled while partial breaking of $\mathcal{N} = 2$ supersymmetry

is realized as before. It shows that the fermionic shift symmetry is due to the free NG fermion.

In [9], we provide a manifestly $\mathcal{N} = 2$ supersymmetric formulation of the $\mathcal{N} = 2$ $U(N)$ gauge model with/without $\mathcal{N} = 2$ hypermultiplets. It is convenient to use $\mathcal{N} = 2$ superfields in harmonic superspace: vector multiplet $V^{++a}(x^m, \theta^+, \bar{\theta}^+) \ni (A^a, v_m^a, \lambda_\alpha^{aI}, D^{aIJ})$ and hypermultiplet q^{+a} in the adjoint representation. The action for the gauge field and the $U(N)$ gauged matter is

$$S_{V+q} = -\frac{i}{4} \int d^4x [(D)^4 \mathcal{F}(W) - (\bar{D})^4 \bar{\mathcal{F}}(\bar{W})] - \int dud\zeta \zeta^{(-4)} \tilde{q}_a^+ \mathcal{D}^{++} q^{+a} \quad (12)$$

where W is the curvature of V^{++} and $\mathcal{D}^{++} q^{+a} = D^{++} q^{+a} + iV^{++c} i f_{cb}^a q^{+b}$. The electric FI term, $S_e = \int dud\zeta \zeta^{(-4)} \text{tr}(\Xi^{++} V^{++}) + c.c.$, shifts the dual auxiliary field D_D^{aIJ} in $W_D^a \equiv \mathcal{F}_a$ by an imaginary constant. So we introduce the magnetic FI term S_m so as to shift the auxiliary field D^{aIJ} as $D^{aIJ} \rightarrow \mathbf{D}^{aIJ} = D^{aIJ} + 4i\xi^{IJ} \delta_0^a$. By this, the $\mathcal{N} = 2$ supersymmetry transformation law changes to $\delta_\eta \lambda^{aI} = (\mathbf{D}^a)^I_J \eta^J + \dots$, under which the total action $S = S_V + S_q^{\text{gauged}} + S_e + S_m$ is invariant. We find that on the Coulomb phase the $\mathcal{N} = 2$ supersymmetry is partially broken to $\mathcal{N} = 1$ spontaneously, and (8) is reproduced by fixing $SU(2)$ appropriately. A generalization to the case with $\mathcal{N} = 2$ local supersymmetry is discussed in [13].

REFERENCES

1. J. T. Lopuszanski, Rept. Math. Phys. **13** (1978) 37.
2. I. Antoniadis, H. Partouche and T. R. Taylor, Phys. Lett. B **372** (1996) 83 [arXiv:hep-th/9512006].
3. S. Ferrara, L. Girardello and M. Porrati, Theories,” Phys. Lett. B **376** (1996) 275 [arXiv:hep-th/9512180].
4. K. Fujiwara, H. Itoyama and M. Sakaguchi, Prog. Theor. Phys. **133** (2005) 429 [arXiv:hep-th/0409060]; in the proceedings of SUSY2004 [arXiv:hep-th/0410132].
5. K. Fujiwara, H. Itoyama and M. Sakaguchi, Nucl. Phys. B **723** (2005) 33 [arXiv:hep-th/0503113].
6. K. Fujiwara, arXiv:hep-th/0609039.
7. L. Andrianopoli, M. Bertolini, A. Ceresole, R. D’Auria, S. Ferrara and P. Fre’, Nucl. Phys. B **476** (1996) 397 [arXiv:hep-th/9603004]; L. Andrianopoli, M. Bertolini, A. Ceresole, R. D’Auria, S. Ferrara, P. Fre and T. Magri, J. Geom. Phys. **23** (1997) 111 [arXiv:hep-th/9605032].
8. B. de Wit, J. W. van Holten and A. Van Proeyen, Nucl. Phys. B **167** (1980) 186; J. Bagger and E. Witten, Nucl. Phys. B **222** (1983) 1; B. de Wit, P. G. Lauwers, R. Philippe, S. Q. Su and A. Van Proeyen, Phys. Lett. B **134** (1984) 37; B. de Wit and A. Van Proeyen, Nucl. Phys. B **245** (1984) 89; B. de Wit, P. G. Lauwers and A. Van Proeyen, Nucl. Phys. B **255** (1985) 569; E. Cremmer, C. Kounnas, A. Van Proeyen, J. P. Derendinger, S. Ferrara, B. de Wit and L. Girardello, Nucl. Phys. B **250** (1985) 385; H. Itoyama, L. D. McLerran, T. R. Taylor and J. J. van der Bij, Nucl. Phys. B **279** (1987) 380; R. D’Auria, S. Ferrara and P. Fre, Nucl. Phys. B **359** (1991) 705.
9. K. Fujiwara, H. Itoyama and M. Sakaguchi, Nucl. Phys. B **740** (2006) 58 [arXiv:hep-th/0510255]; in the proceedings of “Frontier of Quantum Physics” [arXiv:hep-th/0602267].
10. A. S. Galperin, E. A. Ivanov, V. I. Ogievetsky and E. S. Sokatchev, “Harmonic superspace,” Cambridge University Press, 2001.
11. F. Cachazo, M. R. Douglas, N. Seiberg and E. Witten, JHEP **0212** (2002) 071 [arXiv:hep-th/0211170].
12. S. Ferrara and B. Zumino, Nucl. Phys. B **87**, 207 (1975); H. Itoyama, M. Koike and H. Takashino, Mod. Phys. Lett. A **13**, 1063 (1998) [arXiv:hep-th/9610228].
13. H. Itoyama and K. Maruyoshi, arXiv:hep-th/0603180; K. Maruyoshi, arXiv:hep-th/0607047.